



**MUSCLE
GENERALITY**
DANIL HAMMOUDI.MD


- We have over 600 major muscles.
- We have 240 muscles that have specific jobs

Sartorius muscle
It is the longest muscle in the body. It runs from the anterior superior iliac spine of the hip bone to the medial surface of the shaft of the tibia.




http://www.brookscole.com/chemistry_d/templates/student_resources/shared_resources/animations/muscles/muscles.html






Function of Muscles

- Produce movement
- Maintain posture
- Stabilize joints
- Support soft tissue
- Guard openings to the internal body
- Generate heat



Muscle Function

- Skeletal muscles are responsible for all locomotion
- Cardiac muscle is responsible for coursing the blood through the body
- Smooth muscle helps maintain blood pressure, and squeezes or propels substances (i.e., food, feces) through organs
- Muscles also maintain posture, stabilize joints, and generate heat



1. Production of Movement

- Movement of body parts and of the environment
- Movement of blood through the heart and the circulatory vessels.
- Movement of lymph through the lymphatic vessels
- Movement of food (and, subsequently, food waste) through the GI tract
- Movement of bile out of the gallbladder and into the digestive tract
- Movement of urine through the urinary tract
- Movement of semen through the male reproductive tract and female reproductive tract
- Movement of a newborn through the birth canal

2. Maintenance of posture

- Muscle contraction is constantly allowing us to remain upright.
- The muscles of your neck are keeping your head up right now.
- As you stand, your leg muscles keep you on two feet.

3. Thermogenesis

- Generation of heat. Occurs via shivering – an involuntary contraction of skeletal muscle.

Agonist

A muscle that causes motion.

Antagonist

A muscle that can move the joint opposite to the movement produced by the agonist.

Target

The primary muscle intended for exercise.

Synergist

A muscle that assists another muscle to accomplish a movement.

Stabilizer


A muscle that contracts with no significant movement

4. Stabilization of joints

- Muscles keep the tendons that cross the joint nice and taut. This does a wonderful job of maintaining the integrity of the joint.

Types of Muscles

- Skeletal muscle
- Cardiac muscle
- Smooth muscle



THREE HISTOLOGICAL TYPES OF MUSCLE FIBERS

- 1) Skeletal (Striated) = striated (striped in appearance under microscope), voluntary (under conscious control)
- 2) Cardiac = striated, involuntary
- 3) Smooth = non-striated, involuntary

Origin (b): muscle attachment that moves least, generally more proximal.

Insertion (a): muscle attachment that moves most, generally more distal.

Abduction: Lateral movement away from the midline of the body

Adduction: Medial movement toward the midline of the body

Circumduction: circular movement (combining flexion, extension, adduction, and abduction) with no shaft rotation

Extension: Straightening the joint resulting in an increase of angle

Eversion: Moving sole of foot away from medial plane

Flexion: Bending the joint resulting in a decrease of angle

Hyperextension: extending the joint beyond anatomical position

Inversion: Moving sole of foot toward medial plane

Pronation: Internal rotation resulting in appendage facing downward

Protrusion: Moving anteriorly (eg: chin out)

Supination: External rotation resulting in appendage facing upward

Retrusion: Moving posteriorly (eg: chin in)

Rotation: Rotary movement around the longitudinal axis of the bone

Muscle Types

- Smooth
 - blood vessels
 - autonomic
- Striated
 - voluntary
 - skeletal
- Cardiac
 - network
 - rhythmic

The three types of human muscle tissue

- **Smooth, nonstriated muscle** is found in the walls of the hollow viscera and blood vessels.
- **Skeletal, striated muscle** is attached to the skeleton and provides the force for movement of the bony leverage system.
- **Cardiac, striated muscle** is found only in the heart.

Muscle fiber types are classified by

- Anatomical appearance: Anatomical appearance: red versus white red versus white
- Muscle function: Muscle function: fast fast-slow or fatigable versus slow or fatigable versus fatigue resistant fatigue resistant
- Biochemical properties: Biochemical properties: such as high or low such as high or low aerobic capacity aerobic capacity
- Histochemical Histochemical properties: properties: such as enzyme such as enzyme profile

Muscle Classification

- As many as eight types
- Red (type I)
 - long term
 - slow contractions
- White (type IIa)
 - short term
 - fast contractions
- White (type IIb)
- No change from one type to another
 - change within fast types

Muscle Similarities

- Skeletal and smooth muscle cells are elongated and are called muscle fibers
- Muscle contraction depends on two kinds of myofilaments – actin and myosin
- Muscle terminology is similar
 - Sarcolemma – muscle plasma membrane
 - Sarcoplasm – cytoplasm of a muscle cell
 - Prefixes – myo, mys, and sarco all refer to muscle

Muscle Classification cont...

- Change in the nerve root supply will change the muscles twitch properties.
- No gender differences.
- No change in the relative % of each type with training.
- Your birth determines your activity?

Functional Characteristics of Muscle Tissue

- Excitability, or irritability – the ability to receive and respond to stimuli
- Contractility – the ability to shorten forcibly
- Extensibility – the ability to be stretched or extended
- Elasticity – the ability to recoil and resume the original resting length

Interactions of Skeletal Muscles

- Skeletal muscles work together or in opposition
- Muscles only pull (never push)
- As muscles shorten, the insertion generally moves toward the origin
- Whatever a muscle (or group of muscles) does, another muscle (or group) “undoes”

Bone-Muscle Relationships: Lever Systems




Figure 10.2b

Bone-Muscle Relationships: Lever Systems

- Lever – a rigid bar that moves on a fulcrum, or fixed point
- Effort – force applied to a lever
- Load – resistance moved by the effort

Lever Systems: Classes

- First class – the fulcrum is between the load and the effort
- Second class – the load is between the fulcrum and the effort
- Third class – the effort is applied between the fulcrum and the load

Bone-Muscle Relationships: Lever Systems

Effort \times length of effort arm = load \times length of load arm
(force \times distance) = (resistance \times distance)




Figure 10.2a

Lever Systems: First Class





Figure 10.3a

Lever Systems: Second Class




(b) Second-class lever


Figure 10.3b

Skeletal Muscle – the organ

- Skeletal muscle organs are dominated by muscle tissue but also contain nervous, vascular and assorted connective tissues.
- The whole muscle is surrounded by a layer of dense irregular connective tissue known as the **epimysium** (*epi*=?, *mysium*=muscle).



Lever Systems: Third Class




(c) Third-class lever

Figure 10.3c

- Epimysium surrounds several bundles known as **fascicles**.
- Each fascicle is a bundle of super-long skeletal muscle cells (**muscle fibers**), surrounded by a layer of dense irregular CT called the **perimysium** (*peri*=around).


Skeletal Muscle – the organ



- Each muscle cell extends the length of the whole muscle organ and is surrounded by a fine layer of loose connective tissue, the **endomysium**.
- The epi-, peri-, and endomysium are all continuous with one another.



Muscle Histology

Three Muscle Types















Structure and Organization of Skeletal Muscle

STRUCTURE AND ORGANIZATIONAL LEVEL	DESCRIPTION	CONNECTIVE TISSUE WRAPPINGS
<p>Muscle (organ)</p> <p>Epimysium Fascicle Muscle Tendon</p>	Consists of hundreds to thousands of muscle cells, plus connective tissue wrappings, blood vessels, and nerve fibers.	Covered externally by the epimysium
<p>Fascicle (a portion of the muscle)</p> <p>Muscle fiber (cell) Perimysium</p> <p>Part of a fascicle</p>	Discrete bundle of muscle cells, segregated from the rest of the muscle by a connective tissue sheath.	Surrounded by a perimysium
<p>Muscle fiber (cell)</p> <p>Nucleus Sarcolemma surrounded by endomysium Myofibril</p> <p>Part of a muscle fiber Striations</p>	Elongated multinucleate cell; has a banded (striated) appearance.	Surrounded by the endomysium



Table 9.1a








1. Motor end plates
2. Terminal axon fibers
3. Skeletal muscle fibers




Skeletal muscles (C) are stimulated by nerve impulses carried by motor neurons.

The axon (A) that carries the impulse away from the nerve body ends on muscle fibers in little pads called **motor end plates** (B).




Myotendinous Junction




Smooth muscle cells are spindle shaped and uninucleate. (B).

Locations: walls of hollow organs, i.e. stomach, intestine, uterus, ureter



Functions: involuntary movement - i.e. churning of food, movement of urine from the kidney to the bladder, partuition





Smooth muscle - with cells more separated so as to see their extent and shape better, and the central position of their nuclei.
A loose, irregular connective tissue (endomysium) lies between the cells.
Nuclei seen in this c.t. belong to fibroblasts mainly.









Smooth muscle with wrinkled nuclei due to contraction of cells.







Cross-cut of skeletal muscle to show connective tissue partitioning of muscle into groups or bundles of fibers. Endomysium is very delicate and lies between individual fibers, while perimysium is more visible and lies around a group of fibers. Epimysium is not seen here but ensheaths a whole muscle. In this picture notice the presence of small blood vessels in both perimysium and endomysium. Notice also the cross-cuts of myofibrils within the muscle cells, making them look grainy.



Longitudinal view of skeletal muscle cell with unusually clear cross-striations. This muscle is stretched, so that the A band is widely split.
a) Z disc
b) A band, split -- with pale H band in the middle
c) the line lies right in an H band
d) width of I band, with Z disc in the middle
e) pointing to a practically invisible thin line, the sarcolemma (or cell membrane), which lies outside the pale peripheral nucleus seen to the right.






Skeletal Muscle – Blood & Nerve Supply

- Each skeletal muscle is typically supplied by one nerve, an artery and one or more veins.
 - What is the function of each of these 3 items?
- They all enter/exit via the connective tissue coverings and branch extensively.


tibial nerve

Skeletal Muscle Attachments

- Most span joints and are attached to bones.
 - The attachment of the muscle to the immovable bone in a joint is its **origin**, while the attachment to the moveable bone is its **insertion**.



Skeletal Muscle Microanatomy




- Each skeletal muscle cell is known as a skeletal muscle fiber because they are so long.
 - Their diameter can be up to 100µm and their length can be as long as 30cm.
 - They're so large because a single skeletal muscle cell results from the fusion of hundreds of embryonic precursor cells called **myoblasts**.
 - A cell made from the fusion of many others is known as a **syncytium**.
 - Each skeletal muscle fiber will have multiple nuclei. Why?

Muscle attachments may be direct or indirect.


Direct attachments are less common. The epimysium is fused to a periosteum or a perichondrium.



Indirect attachments are typical. The muscle CT extends and forms either a cordlike structure (a tendon) or a sheetlike structure (aponeurosis) which attaches to the periosteum or perichondrium.




Muscle fiber PM is known as **sarcolemma**
Muscle fiber cytoplasm is known as **sarcoplasm**




Sarcolemma has invaginations that penetrate through the cell called **transverse tubules** or **T tubules**.

Sarcoplasm has lots of mitochondria (*why?*), lots of glycogen granules (to provide glucose for energy needs) as well as **myofibrils** and **sarcoplasmic reticuli**.




Sarcoplasmic Reticulum

- Muscle cell version of the smooth endoplasmic reticulum.
- Functions as a calcium storage depot in muscle cells.
- Loose network of this membrane bound organelle surrounds all the myofibrils in a muscle fiber. We will see why this is so important soon.




Myofibrils

- Each muscle fiber contains rodlike structures called myofibrils that extend the length of the cell. They are basically long bundles of protein structures called myofilaments and their actions give muscle the ability to contract.
- The myofilaments are classified as thick filaments and thin filaments.



Myofibers

- **Myofiber**= one muscle cell= muscle fiber
- Myofiber is syncytial (multinucleate).
- Myofibers are long!



Myofibers

- In order of decreasing size...
- **Myofiber**= entire cell.
- **Myofibrils**: bundles of myofilaments inside myofiber.
- **Myofilaments**: actin and myosin proteins.


•Myofibers

Myofibrils


- Myofibrils are densely packed, rodlike contractile elements
- They make up most of the muscle volume
- The arrangement of myofibrils within a fiber is such that a perfectly aligned repeating series of dark A bands and light I bands is evident

Myofibrils

- Each myofibril is made up 1000's of repeating individual units known as **sarcomeres** (pictured below)
- Each sarcomere is an ordered arrangement of thick and thin filaments. Notice that it has:
 - regions of thin filaments by themselves (pinkish fibers)
 - a region of thick filaments by themselves (purple fibers)
 - regions of thick filaments and thin filaments overlapping.



Myofibrils




PLAY InterActive Physiology ©:
Anatomy Review: Skeletal Muscle Tissue, pages 7-8

Figure 9.3b

The portion of the sarcomere which does not contain any thick filament is known as the **I band**. The I band contains only thin filament and is light under the microscope (*it is isotropic*).

- One I band is actually part of 2 sarcomeres at once.




In the middle of the H zone is a structure called the **M line** which functions to hold the thick filaments to one another

“SARCOTERMS”


- **Sarcolemma**: pm of myofiber
- **Sarcoplasm**: cytoplasm of myofiber
- **Sarcoplasmic reticulum (SR)**: ER of myofiber
- **Sarcomere**: contractile unit inside myofiber

Here we have several different cross sections of a myofibril. Why are they different?




Sarcomere

- The sarcomere is flanked by 2 protein structures known as **Z discs**.
- The portion of the sarcomere which contains the thick filament is known as the **A band**. A stands for *anisotropic* which is a fancy way of saying that it appears dark under the microscope.
 - The A band contains a **zone of overlap** (btwn thick & thin filaments) and an **H zone** which contains only thick filaments



Sarcomeres


- The smallest contractile unit of a muscle
- The region of a myofibril between two successive Z discs
- Composed of myofilaments made up of contractile proteins
 - Myofilaments are of two types
 - thick and thin



Myofilaments: Banding Pattern

- Thin filaments do not overlap thick filaments in the lighter H zone
- M lines appear darker due to the presence of the protein desmin

Sarcomeres



PLAY InterActive Physiology ©:
Anatomy Review: Skeletal Muscle Tissue, page 9

Figure 9.3c

Myofilaments: Banding Pattern




Figure 9.3c,d

Myofilaments: Banding Pattern




- Thick filaments – extend the entire length of an A band
- Thin filaments – extend across the I band and partway into the A band
- Z-disc – coin-shaped sheet of proteins (connectins) that anchors the thin filaments and connects myofibrils to one another

Structure and Organization of

TABLE 9.1 Structure and Organizational Levels of Skeletal Muscle

STRUCTURE AND ORGANIZATIONAL LEVEL	DESCRIPTION	CONNECTIVE TISSUE WRAPPINGS
Myofibril or fibril (complex organelle composed of bundles of myofilaments)	Rodlike contractile element; myofibrils occupy most of the muscle cell volume; composed of sarcomeres arranged end to end; appear banded, and bands of adjacent myofibrils are aligned	
Sarcomere (a segment of a myofibril)	The contractile unit, composed of myofilaments made up of contractile proteins	
Myofilament or filament (extended macromolecular structure)	Contractile myofilaments are of two types—thick and thin; the thick filaments contain bundled myosin molecules; the thin filaments contain actin molecules (plus other proteins); the sliding of the thin filaments past the thick filaments produces muscle shortening. Elastic filaments (not shown here) maintain the organization of the A band and provide for elastic recoil when muscle contraction ends	


Table 9.1b



Muscle Contraction: The Sliding Filament Hypothesis

- Place your right palm on the back of your left hand. Now slide your right palm toward your left elbow.
 - What happened to the distance between your elbows?
 - It got shorter!
 - This is how muscle contraction occurs.
 - The thin filaments slide over the thick filaments. This pulls the Z discs closer together. When all the sarcomeres in a fiber do this, the entire fiber gets shorter which pulls on the endomysium, perimysium, epimysium and attached tendon and then pulls on the b.

% Tension Developed




Here is what happens as the filaments slide and the sarcomere and the muscle fiber shortens.

In the process of contraction, what happens to the:

- Distance btwn Z discs
- Length of the A band
- Length of the H zone
- Length of the I band


Here are 2 electron micrographs of the same sarcomere. Do you see the Z discs, A band, H zone, M line, and I bands? How do the 2 pictures differ? What happened?



Ultrastructure of Myofilaments:




(a) Myosin molecule



(b) Portion of a thick filament

Figure 9.4a,b



Ultrastructure of Myofilaments: Thin Filaments

- Thin filaments are chiefly composed of the protein actin
- Each actin molecule is a helical polymer of globular subunits called G actin
- The subunits contain the active sites to which myosin heads attach during contraction
- Tropomyosin and troponin are regulatory subunits bound to actin

Ultrastructure of Myofilaments: Thick Filaments


- Thick filaments are composed of the protein myosin
- Each myosin molecule has a rod-like tail and two globular heads
 - Tails – two interwoven, heavy polypeptide chains
 - Heads – two smaller, light polypeptide chains called cross bridges

Myofilaments


- 2 types of myofilaments (**thick & thin**) make up myofibrils.
- Thick myofilaments are made the protein **myosin**

A single myosin protein resembles 2 golf clubs whose shafts have been twisted about one another

About 300 of these myosin molecules are joined together to form a single thick filament




- Each thin filament is made up of 3 different types of protein: **actin, tropomyosin, and troponin**.
 - Each thin filament consists of a long helical double strand. This strand is a polymer that resembles a string of beads. Each “bead” is the globular protein **actin**. On each actin subunit, there is a **myosin binding site**.
 - Loosely wrapped around the actin helix and covering the myosin binding site is the filamentous protein, **tropomyosin**.
 - Bound to both the actin and the tropomyosin is a trio of proteins collectively known as **troponin**.



Arrangement of the Filaments in a Sarcomere



- Longitudinal section within one sarcomere



(d) Longitudinal section of filaments within one sarcomere of a myofibril


Figure 9.4d

Note the relationship between the thin and thick filaments



Actin Myosin Crossbridge 3D Animation

Ultrastructure of Myofilaments: Thin Filaments



(c) Portion of a thin filament

Figure 9.4c

Sarcoplasmic Reticulum (SR)


- SR is an elaborate, smooth endoplasmic reticulum that mostly runs longitudinally and surrounds each myofibril
- Paired terminal cisternae form perpendicular cross channels
- Functions in the regulation of intracellular calcium levels

Sarcoplasmic Reticulum (SR)

- Elongated tubes called T tubules penetrate into the cell's interior at each A band-I band junction
- T tubules associate with the paired terminal cisternae to form triads

T-Tubules and the SR

- Each muscle fiber has many T-tubules
 - Typically each myofibril has a branch of a T-tubule encircling it at each A-I junction
- At each A-I junction, the SR will expand and form a dilated sac (terminal cisterna).



Each T-tubule will be flanked by a terminal cisterna. This forms a so-called triad consisting of 2 terminal cisternae and one T-tubule branch.

Sarcoplasmic Reticulum (SR)




Figure 9.5

Triad Relationships

- T tubules and SR provide tightly linked signals for muscle contraction
- A double zipper of integral membrane proteins protrudes into the intermembrane space
- T tubule proteins act as voltage sensors
- SR foot proteins are receptors that regulate Ca^{2+} release from the SR cisternae

PLAY InterActive Physiology ©:
Anatomy Review: Skeletal Muscle Tissue, page 10

T Tubules

- T tubules are continuous with the sarcolemma
- They conduct impulses to the deepest regions of the muscle
- These impulses signal for the release of Ca^{2+} from adjacent terminal cisternae

Sliding Filament Model of Contraction

- Thin filaments slide past the thick ones so that the actin and myosin filaments overlap to a greater degree
- In the relaxed state, thin and thick filaments overlap only slightly
- Upon stimulation, myosin heads bind to actin and sliding begins

SKELETAL MUSCLE

Skeletal Muscle

- The three connective tissue sheaths are:
 - Endomysium – fine sheath of connective tissue composed of reticular fibers surrounding each muscle fiber
 - Perimysium – fibrous connective tissue that surrounds groups of muscle fibers called fascicles
 - Epimysium – an overcoat of dense regular connective tissue that surrounds the entire muscle


Skeletal muscles are composed of fibers (cells) bound together by connective tissue.

- Connective tissue associated with muscle [endomysium, perimysium, epimysium, fascia]
- Muscle attachments
- Tendons are bands of dense connective tissue attaching muscle to bone.

- Most moveable attachment is **insertion**
- Least moveable attachment is **origin**


- **Aponeurosis** is a flattened sheet like muscle attachment.
- Four principal fiber patterns in skeletal muscle
 1. Parallel fibers [rectus abdominus]
 2. Convergent fibers [pectoralis major]
 3. Circular fibers [orbicularis oris]
 4. Pennate fibers [rectus femoris]



- Muscle cells contract when stimulated by nerve impulses • Isotonic & Isometric Contractions
- Motor Unit is a motor neuron + muscle fibers it innervates
- Myoneural Junction is the contact site between the end of the motor neuron and muscle fiber
- Motor End Plate is the portion of the sarcolemma in contact with motor neuron.



SKELETAL MUSCLE

- I. CELLS (FIBERS)
 - 1) Very long compared with most other cells, up to several cm long, 10-100 micrometers in diameter
 - 2) Multinucleate, nuclei are located peripherally
 - 3) Development: Mesenchymal cell → Myoblast (proliferative) → Myotubule → Muscle Cell
- II. ARRANGEMENT OF FIBERS - similar to tendon arrangement
 - Blood vessels, lymph vessels, and nerves penetrate muscle with perimysium
 - Endomysium contains capillaries and nerve fibers
- III. STRIATION ULTRASTRUCTURE (Fibers → Myofibrils → Myofilaments)
 - Proteins are *actin* (thin filaments) and *myosin* (thick filaments), also tropomyosin and troponin are associated with thin filaments





Skeletal Muscle

- Skeletal muscle is attached to bone on each end by tendons.
 - Insertion:
 - More movable attachment.
 - Origin:
 - Are pulled towards it.
- Surrounded by connective tissues (around each muscle, subsets, and muscle cell).

Skeletal Muscle



- Motor end plate= sarcolemma at neuromuscular junction

Skeletal Muscle: Attachments

- Muscles attach:
 - Directly – epimysium of the muscle is fused to the periosteum of a bone
 - Indirectly – connective tissue wrappings extend beyond the muscle as a tendon or aponeurosis

Skeletal Muscle

- Motor unit= myofibers innervated by same motor neuron
- Muscles:
 - Graded contractions by how many motor units contract.



Microscopic Anatomy of a Skeletal Muscle Fiber

- Each fiber is a long, cylindrical cell with multiple nuclei just beneath the sarcolemma
- Each cell is a syncytium produced by fusion of embryonic cells


Muscles

- **Twitch** = muscle contraction
- **Summation:**
 - If second stimulus is administered before complete relaxation of muscle.
- **Complete tetanus:**
 - Fusion frequency of stimulation.
 - No visible relaxation between twitches.
 - Smooth sustained contraction.

Microscopic Anatomy of a Skeletal Muscle Fiber

- Sarcoplasm has numerous glycosomes and a unique oxygen-binding protein called myoglobin
- Fibers contain the usual organelles, myofibrils, sarcoplasmic reticulum, and T tubules

Muscles



Muscles

• Treppe:

- Second stimulus elicits a stronger response
 - Perhaps due to increase in intracellular Ca^{2+} .

Metabolism

- During some exercise: ATP used faster than can be renewed through cell respiration.
- So: **Phosphocreatine reservoir!**
- $ADP + \text{phosphocreatine} \rightarrow ATP + \text{creatine}$

Contractions

- **Isotonic** contractions:
 - Force of contraction remains constant throughout the shortening process.
- **Isometric** contractions:
 - Length of muscle fibers remain constant.
- **Eccentric** contractions:
 - Force exerted on a muscle to stretch is greater than the force of muscle contraction.
 - Running downhill
 -

Skeletal muscle

- Contraction speed:
 - Slow-twitch.
 - Fast-twitch.
- due to different myosins.
- Red/white meat...!

Tendons

- **Tendons**:
 - Have elasticity.
 - Display recoil.
 - Spring back to resting length.

Individual Fiber Types

Fast fibers

• Type IIb

- Fast Fast-twitch fibers twitch fibers
- Fast Fast-glycolytic glycolytic fibers fibers

• Type IIa

- Intermediate fibers Intermediate fibers
- Fast Fast-oxidative oxidative glycolytic glycolytic fibers

Slow fibers

• Type I fibers

- Slow Slow-twitch fibers twitch fibers
- Slow Slow-oxidative oxidative fibers

Myofibers

- **Slow-twitch:**
 - Red fibers.
 - High oxidative capacity for aerobic respiration.
 - Many: mitochondria, capillaries
 - Myoglobin (like hemoglobin) for oxygen.
 - Postural muscles


Muscle Spindle Apparatus

- **Muscle spindle apparatus**
 - Length detector.
 - Contains thin muscle cells called intrafusal fibers.
 - Reflex contraction in response to rapid stretch.
 - Stimulated by γ motor neurons from spinal cord.
 - Helps maintain muscle tone (resting muscle length and state of tension).
 - Extrafusal fibers (rest of muscle!): stimulated by α motor neurons from spinal cord.

Myofibers

- **Fast-twitch (type IIX fibers):**
 - White fibers.
 - respire anaerobically.
 - much glycogen.
- **Fast-twitch oxidative (type IIA fibers):**
 - Also white fibers.
- People vary genetically in proportion of fast- and slow-twitch fibers in their muscles.

Spindle apparatus




Exercise

- Weight lifting: hypertrophy.
- Endurance training: more mitochondria.

Golgi Tendon Organ

•Golgi tendon organ


- Helps prevent excessive muscle contraction or excessive passive muscle stretching.
- A reflex.




Skeletal Muscles

- **Agonist** muscle:
 - Prime mover.
- **Antagonist** muscle:
 - Flexors and extensors that act on the same joint to produce opposite actions.

CARDIAC MUSCLE



Cardiac Muscle Fiber




Intercalated Disc
Nucleus


Reciprocal Innervation

- **Reciprocal innervation:**
- motor neurons of antagonistic muscles inhibit each other (through interneurons) so they don't both contract simultaneously.
- When limb is flexed, antagonistic extensor muscles are passively stretched.

Cardiac Muscle




Gap junctions Myocardial cells




CARDIAC MUSCLE

INTERCALATED DISCS


Intercalated Discs



Cardiac Muscle Fibers (transverse section)




INTERCALATED DISCS





• SMOOTH MUSCLE

PURKINJE FIBERS


Purkinje Fibers
(specialized
conducting
tissue)



Purkinje Fibers (trans. section)



LOCATIONS OF SMOOTH MUSCLE



DIGESTIVE TRACT
 DUCTS OF GLANDS
 RESPIRATORY PASSAGES
 URINARY & GENITAL TRACT
 ARTERIES AND VEINS
 PILIERECTOR MUSCLES
 IRIS & CILIARY BODY
 UTERUS
 BLADDER
 STOMACH


Cardiac Muscle

- Like skeletal muscle:
 - - striated: actin and myosin in sarcomeres.
 - - contract via sliding-filament mechanism.
- Unique to cardiac muscle:
 - - Adjacent myocardial cells joined by gap junctions= intercalated discs=electrical synapse.


Smooth Muscle

- Not striated.
- NO sarcomeres.
- Lots of actin, some myosin
- Can contract even when very stretched.
- Graded contractions

Smooth Muscle



Myofiber anatomy



Smooth Muscle


- Actin all over the cell, linked by myosin (web-like pattern, not striations).
- Rise in Ca^{2+} -> Ca^{2+} binds with **calmodulin** -> activates **MLCK** (a kinase) -> Myosin heads are phosphorylated and can bind to actin.

The Overload Principle

- Forcing a muscle to work promotes increased muscular strength
- Muscles adapt to increased demands
- Muscles must be overloaded to produce further gains

Myofiber anatomy

- Sarcoplasmic reticulum lies next to T tubules.
- Transverse tubules = infoldings of sarcolemma.



Smooth Muscle

- Composed of spindle-shaped fibers with a diameter of 2-10 μm and lengths of several hundred μm
- Lack the coarse connective tissue sheaths of skeletal muscle, but have fine endomysium
- Organized into two layers (longitudinal and circular) of closely apposed fibers

Smooth Muscle

- Found in walls of hollow organs (except the heart)
- Have essentially the same contractile mechanisms as skeletal muscle

Innervation of Smooth Muscle

- Smooth muscle lacks neuromuscular junctions
- Innervating nerves have bulbous swellings called varicosities
- Varicosities release neurotransmitters into wide synaptic clefts called diffuse junctions

Smooth Muscle




Figure 9.24

Innervation of Smooth Muscle




Figure 9.25

Peristalsis

- When the longitudinal layer contracts, the organ dilates and contracts
- When the circular layer contracts, the organ elongates
- Peristalsis – alternating contractions and relaxations of smooth muscles that mix and squeeze substances through the lumen of hollow organs

Microscopic Anatomy of Smooth Muscle

- SR is less developed than in skeletal muscle and lacks a specific pattern
- T tubules are absent
- Plasma membranes have pouchlike infoldings called caveoli

Microscopic Anatomy of Smooth Muscle

- Ca^{2+} is sequestered in the extracellular space near the caveoli, allowing rapid influx when channels are opened
- There are no visible striations and no sarcomeres
- Thin and thick filaments are present

Hyperplasia

- Certain smooth muscles can divide and increase their numbers by undergoing hyperplasia
- This is shown by estrogen's effect on the uterus
 - At puberty, estrogen stimulates the synthesis of more smooth muscle, causing the uterus to grow to adult size
 - During pregnancy, estrogen stimulates uterine growth to accommodate the increasing size of the growing fetus

Proportion and Organization of Myofilaments in Smooth Muscle

- Ratio of thick to thin filaments is much lower than in skeletal muscle
- Thick filaments have heads along their entire length
- There is no troponin complex

Types of Smooth Muscle: Single Unit

- The cells of single-unit smooth muscle, commonly called visceral muscle:
 - Contract rhythmically as a unit
 - Are electrically coupled to one another via gap junctions
 - Often exhibit spontaneous action potentials
 - Are arranged in opposing sheets and exhibit stress-relaxation response

Proportion and Organization of Myofilaments in Smooth Muscle

- Thick and thin filaments are arranged diagonally, causing smooth muscle to contract in a corkscrew manner
- Noncontractile intermediate filament bundles attach to dense bodies (analogous to Z discs) at regular intervals

Types of Smooth Muscle: Multiunit

- Multiunit smooth muscles are found:
 - In large airways to the lungs
 - In large arteries
 - In arrector pili muscles
 - Attached to hair follicles
 - In the internal eye muscles

Types of Smooth Muscle: Multiunit

- Their characteristics include:
 - Rare gap junctions
 - Infrequent spontaneous depolarizations
 - Structurally independent muscle fibers
 - A rich nerve supply, which, with a number of muscle fibers, forms motor units
 - Graded contractions in response to neural stimuli

Muscle types







Muscle type	Activity
Skeletal muscle 	 Strong, quick discontinuous voluntary contraction
Cardiac muscle 	 Strong, quick continuous involuntary contraction
Smooth muscle 	 Weak, slow involuntary contraction

TABLE 9.3 Comparison of Skeletal, Cardiac, and Smooth Muscle

CHARACTERISTIC	SKELETAL	CARDIAC	SMOOTH
Body location	Attached to bones or (some facial muscles) to skin	Walls of the heart	Single-unit muscle in walls of hollow visceral organs (other than the heart); multiunit muscle in intrinsic eye muscles, airways, large arteries
Cell shape and appearance	Single, very long, cylindrical, multinucleate cells with obvious striations	Branching chains of cells; uni- or binucleate; striations	Single, fusiform, uninucleate; no striations




Table 9.3.1

TABLE 9.3 Comparison of Skeletal, Cardiac, and Smooth Muscle

CHARACTERISTIC	SKELETAL	CARDIAC	SMOOTH
Connective tissue components	Epimysium, perimysium, and endomysium	Endomysium attached to fibrous skeleton of heart	Endomysium
Presence of myofibrils composed of sarcomeres	Yes	Yes, but myofibrils are of irregular thickness	No, but actin and myosin filaments are present throughout; dense bodies anchor actin filaments
Presence of T tubules and site of invagination	Yes; two in each sarcomere at A-I junctions	Yes; one in each sarcomere at Z disc; larger diameter than those of skeletal muscle	No; only caveolae
Elaborate sarcoplasmic reticulum	Yes	Less than skeletal muscle (1-8% of cell volume); scant terminal cisternae	Equivalent to cardiac muscle (1-8% of cell volume); some SR contacts the sarcolemma




Table 9.3.2

Comparison of Skeletal, Cardiac and Smooth Muscle Cells

Skeletal Muscle Cell:	Cardiac Muscle:	Smooth Muscle Cell:
Elongated Cells	Branching Cells	Spindle-Shaped Cell
Multiple Peripheral Nuclei	Single Central Nucleus	Single Central Nucleus
Visible Striations	Visible Striations	Lack Visible Striations
Voluntary	Involuntary	Involuntary




Table 9.3.3

TABLE 9.3 Comparison of Skeletal, Cardiac, and Smooth Muscle

CHARACTERISTIC	SKELETAL	CARDIAC	SMOOTH
Presence of gap junctions	No	Yes; at intercalated discs	Yes; in single-unit muscle
Cells exhibit individual neuromuscular junctions	Yes	No	Not in single-unit muscle; yes in multiunit muscle
Regulation of contraction	Voluntary via axon terminals of the somatic nervous system	Involuntary; intrinsic system regulation; also autonomic nervous system controls; hormones, stretch	Involuntary; autonomic nerves, hormones, local chemicals; stretch
Source of Ca ²⁺ for calcium pulse	Sarcoplasmic reticulum (SR)	SR and from extracellular fluid	SR and from extracellular fluid




Table 9.3.3

TABLE 9.3 Comparison of Skeletal, Cardiac, and Smooth Muscle








CHARACTERISTIC	SKELETAL	CARDIAC	SMOOTH
Site of calcium regulation	Troponin on actin-containing thin filaments 	Troponin on actin-containing thin filaments 	Calmodulin in the sarcoplasm 
Presence of pacemakers(s)	No	Yes	Yes (in single-unit muscle only)
Effect of nervous system stimulation	Excitation	Excitation or inhibition	Excitation or inhibition
Speed of contraction	Slow to fast 	Slow 	Very slow 
Rhythmic contraction	No	Yes	Yes in single-unit muscle
Response to stretch	Contractile strength increases with degree of stretch (to a point)	Contractile strength increases with degree of stretch	Stress-relaxation response
Respiration	Aerobic and anaerobic	Aerobic	Mainly aerobic

Table 9.3.4

Other Important Terms

- **Flaccid paralysis**
 - Weakness or loss of muscle tone typically due to injury or disease of motor neurons
- **Spastic paralysis**
 - Sustained involuntary contraction of muscle(s) with associated loss of function
 - *How do flaccid and spastic paralysis differ?*
- **Spasm**
 - A sudden, involuntary smooth or skeletal muscle twitch. Can be painful. Often caused by chemical imbalances.



Myasthenia Gravis


- My=muscle, asthen=weakness, gravi=heavy
- Autoimmune disease where antibodies attack the ACh receptors on neuromuscular junctions.
- Results in progressive weakening of the skeletal muscles. Why?
- Treated w/ **anticholinesterases** such as neostigmine or physostigmine. These decrease the activity of acetylcholinesterase.
 - *Why would this help someone with myasthenia gravis?*

Other Important Terms

- **Cramp**
 - A prolonged spasm that causes the muscle to become taut and painful.
- **Hypertrophy**
 - Increase in size of a cell, tissue or an organ.
 - In muscles, hypertrophy of the organ is always due to cellular hypertrophy (increase in cell size) rather than cellular **hyperplasia** (increase in cell number)
 - Muscle hypertrophy occurs due to the synthesis of more myofibrils and synthesis of larger myofibrils.

Muscular Dystrophy

- Group of inherited muscle-destroying diseases that generally appear during childhood.
- Dys=faulty; Troph=growth
- Most common is Duchenne muscular dystrophy
 - DMD is caused by an abnormal X-linked recessive gene
 - Diseased muscle fibers lack the protein dystrophin which normally links the cytoskeleton to the ECM and stabilizes the sarcolemma
 - Age of onset is btwn 2 and 10. Muscle weakness progresses. Afflicted individuals usually die of respiratory failure, usually by age 25.



Here is a slide of skeletal muscle from someone with DMD. Look how much connective tissue there is. Lots of adipose tissue too. Why do you think there's so much?

Other Important Terms

- **Atrophy**
 - Reduction in size of a cell, tissue, or organ
 - In muscles, its often caused by disuse. Could a nerve injury result in disuse? Why might astronauts suffer muscle atrophy?
- **Fibrosis**
 - Replacement of normal tissue with heavy fibrous connective tissue (scar tissue). How would fibrosis of skeletal muscles affect muscular strength? How would it affect muscle flexibility?

Muscular Dystrophy

- Muscular dystrophy – group of inherited muscle-destroying diseases where muscles enlarge due to fat and connective tissue deposits, but muscle fibers atrophy

Developmental Aspects

- Muscle tissue develops from embryonic mesoderm called myoblasts
- Multinucleated skeletal muscles form by fusion of myoblasts
- The growth factor *agrin* stimulates the clustering of ACh receptors at newly forming motor end plates

Muscular Dystrophy

- Duchenne muscular dystrophy (DMD)
 - Inherited, sex-linked disease carried by females and expressed in males (1/3500)
 - Diagnosed between the ages of 2-10
 - Victims become clumsy and fall frequently as their muscles fail

Developmental Aspects

- As muscles are brought under the control of the somatic nervous system, the numbers of fast and slow fibers are also determined
- Cardiac and smooth muscle myoblasts do not fuse but develop gap junctions at an early embryonic stage

Muscular Dystrophy

- Progresses from the extremities upward, and victims die of respiratory failure in their 20s
- Caused by a lack of the cytoplasmic protein dystrophin
- There is no cure, but myoblast transfer therapy shows promise

Developmental Aspects: Regeneration

- Cardiac and skeletal muscle become amitotic, but can lengthen and thicken
- Myoblastlike satellite cells show very limited regenerative ability
- Cardiac cells lack satellite cells
- Smooth muscle has good regenerative ability

Developmental Aspects: After Birth

- Muscular development reflects neuromuscular coordination
- Development occurs head-to-toe, and proximal-to-distal
- Peak natural neural control of muscles is achieved by midadolescence
- Athletics and training can improve neuromuscular control

Developmental Aspects: Age Related

- With age, connective tissue increases and muscle fibers decrease
- Muscles become stringier and more sinewy
- By age 80, 50% of muscle mass is lost (sarcopenia)

Developmental Aspects: Male and Female

- There is a biological basis for greater strength in men than in women
- Women's skeletal muscle makes up 36% of their body mass
- Men's skeletal muscle makes up 42% of their body mass

Developmental Aspects: Age Related

- Regular exercise reverses sarcopenia
- Aging of the cardiovascular system affects every organ in the body
- Atherosclerosis may block distal arteries, leading to intermittent claudication and causing severe pain in leg muscles

Developmental Aspects: Male and Female

- These differences are due primarily to the male sex hormone testosterone
- With more muscle mass, men are generally stronger than women
- Body strength per unit muscle mass, however, is the same in both sexes