

EKG The normal and abnormal 2020 UPDATE

D.HAMMOUDI .MD




The Cardiac Cycle

- ▶ **Heart at rest**
 - Blood flows from large veins into atria
 - Passive flow from atria into ventricles
- ▶ **Atria (R & L) contract simultaneously**
 - Blood forced into ventricles
- ▶ **Ventricles (R & L) contract simultaneously**
 - Atrioventricular valves close → “lubb” sound
 - Blood forced into large arteries
- ▶ **Ventricles relax**
 - Semilunar valves close → “dub” sound
- ▶ **Heart at rest**


Depolarization and Impulse Conduction


- ▶ Heart is autorhythmic
- ▶ Depolarization begins in sinoatrial (SA) node
- ▶ Spread through atrial myocardium
- ▶ Delay in atrioventricular (AV) node



Depolarization and Impulse Conduction


- ▶ Spread from atrioventricular (AV) node
 - AV bundle
 - Bundle branches
 - Purkinje fibers





B. Echocardiogram, Apical Four-chamber View

RV	Right ventricle	LV	Left ventricle
RA	Right atrium	LA	Left atrium
x Crux of heart			



S-A NODE

ATRIAL MUSCLE

A-V NODE


COMMON BUNDLE

BUNDLE BRANCHES

PURKINJE FIBERS

VENTRICULAR MUSCLE

ACTION POTENTIALS



Arrhythmia Formation


Arrhythmias can arise from problems in the:

- Sinus node
- Atrial cells
- AV junction
- Ventricular cells




Depolarization and Impulse Conduction



- ▶ Depolarization in SA node precedes depolarization in atria, AV node, ventricles



The ECG Paper

- ▶ Horizontally
 - One small box – 0.04 s
 - One large box – 0.20 s
- ▶ Vertically
 - One large box – 0.5 mV





The ECG Paper (cont)



- ▶ Every 3 seconds (15 large boxes) is marked by a vertical line.
- ▶ This helps when calculating the heart rate.

NOTE: the following strips are not marked but all are 6 seconds long.

Waveforms and Intervals






Fig. 1.4 Relationship between the squares on ECG paper and time. Here, there is one QRS complex per second, so the heart rate is 60 beats/min

■ Q waves

- Can occur normally in several leads
 - Normal Q waves called physiologic
- Physiologic Q waves
 - < .04 sec (40ms)
- Pathologic Q
 - > .04 sec (40ms)

Component	Characteristics
Heart Rate	60-100 bpm
PR Interval	0.12–0.20 sec
QRS Interval	0.06–0.10 sec
QT Interval	Less than half of the R-R interval
ST segment	0.08 sec



P wave (0.08 - 0.10 s) QRS (0.06 - 0.10 s)
P-R interval (0.12 - 0.20 s) Q-T_C interval (≤ 0.44 s)*

$$*QT_C = QT / \sqrt{RR}$$

ECG Manifestations: Electrolyte Imbalance

Diercks DB et al. J Emerg Med. 2004; 27(2); 153-60.

Rosen's EM: Concepts and Clinical Practice, 7th ed, 2010.


Goldberger, AL. Clinical Electrocardiography: A Simplified Approach, 7th ed, 2006.

PR interval	Short	Prolonged
	(Think pre-excitation syndromes such as Wolff-Parkinson-White)	High K Low Ca
QRS duration	Narrow	Wide (>100 msec)
	Low K Low Ca Normal	High K High Ca
QTc interval	Short (<350 msec)	Prolonged (>440 msec)
	High Ca	Low K Low Ca
ST segment	Depressed	Elevated
	Low K High Ca	High K
T wave	Peaked/tall	Flattened
	High K	Low K
U wave	Absent	Present
	Normal	Low K Low Ca
Heart rate	Slow (bradycardia, nodal block)	Fast (tachycardia)
	High K High Ca	Low K Low Ca

	Low	High
Ca	<ul style="list-style-type: none"> • QTc prolonged (hallmark) • U wave • Heart blocks, ventricular dysrhythmias, torsades de pointes 	<ul style="list-style-type: none"> • QTc shortened (hallmark) • ST segment depression and shortening • QRS widening • Rare: bradycardia, bundle branch blocks, high degree AV blocks
K	Early to late findings: <ul style="list-style-type: none"> • T wave: decreased amplitude • T wave: flat or inverted • ST segment depression • U wave • QTc prolonged (at risk for VT or torsades de pointes) 	Early to late findings: <ul style="list-style-type: none"> • T wave: tall, then "peaked" (symmetrical) • P wave flattening • PR interval prolonged • QRS widening • Nodal blocks, escape beats • Sine wave: fusion of QRS and T wave --> VF or asystole

Mg derangements: Nonspecific ECG findings; often co-exist with Ca derangements.
 • Classic teaching: Low Mg level --> QTc prolongation --> torsades de pointes

Waves, intervals and durations on the ECG



The reference level for measuring ST-segment deviation (depression or elevation) is not the TP interval. The correct reference level is the **PR segment**. This level is also called **baseline level** or **isoelectric level**.

EKG Leads

The standard EKG has 12 leads:

3 Standard Limb Leads

3 Augmented Limb Leads


6 Precordial Leads

The axis of a particular lead represents the viewpoint from which it looks at the heart.




Electrocardiogram

- ▶ **P wave**
 - Depolarization of atria
 - Followed by contraction
- ▶ **QRS complex**
 - 3 waves (Q, R, & S)
 - Depolarization of ventricles
 - Followed by contraction
- ▶ **T wave**
 - Repolarization of ventricles




Electrocardiogram


- ▶ P-Q interval
 - Time atria depolarize & remain depolarized
- ▶ Q-T interval
 - Time ventricles depolarize & remain depolarized



Intervals

- **P wave** - atrial depolarization
 - **PR interval** - time from sinoatrial node (S-A) to atrioventricular node (A-Vnode)
 - **QRS Complex** – ventricular depolarization
 - **ST Segment** - beginning of ventricular repolarization
 - **T Wave** - later stages of ventricular repolarization
 - **U Wave** - final component of ventricular repolarization
 - **RR Interval** - represents the time for one complete cardiac cycle
- 

Precordial Leads



Lead Placement

V1 = 4th intercostal space, right border of sternum



V2 = 4th intercostal space, left border of sternum


V3 = midway between V2 and V4

V4 = 5th intercostal space, midclavicular line

V5 = anteroaxillary line at level of V4




V6 = midaxillary line at level of V4 and V5





Lead Placement

- V1 – Right Sternal Border – 4th ICS
- V2 – Left Sternal Border – 4th ICS
- V3 Midway Between V2 and V4
- V4 Midclavicular line – 5th ICS
- V5 Anterior Axillary line – 5th ICS
- V6 Mid axillary line – 5th ICS



- ▶ Lead Placement

- ▶ V1 =

- ▶ V2 =

- ▶ V3 =

- ▶ V4 =

- ▶ V5 =

- ▶ V6 =




- Lead Placement

- V1 = 4th intercostal space, right border of sternum
- V2 = 4th intercostal space, left border of sternum
- V3 = midway between V2 and V4
- V4 = 5th intercostal space, midclavicular line
- V5 = anteroaxillary line at level of V4
- V6 = midaxillary line at level of V4 and V5









RA = Right Arm
LA = Left Arm
RL = Right Leg
LL = Left Leg


RA - White
LA - Black
RL - Green
LL - Red




Electrocardiogram

- ▶ Einthoven's triangle
 - Three standard limb leads
 - Voltage differences between corners of triangle
 - We will use "Lead II"
 - Right shoulder to left leg







-150°
aVR



-30°
aVL




I
0°



III
+120°

aVF
+90°

II
+60°



ECG Recordings: (QRS vector---leftward, inferiorly and posteriorly

3 Bipolar Limb Leads

I = RA vs. LA(+)

II = RA vs. LL(+)



III = LA vs. LL(+)

3 Augmented Limb Leads


aVR = (LA-LL) vs. RA(+)

aVL = (RA-LL) vs. LA(+)

aVF = (RA-LA) vs. LL(+)




6 Precordial (Chest) Leads: Indifferent electrode (RA-LA-LL) vs. chest lead moved from position V_1 through position V_6 .



Normal EKG

- A **positive wave** form (QRS mainly above the baseline) results from the wave of depolarization moving towards the positive end of the lead.
 - e.g.
- A **negative waveform** (QRS mainly below the baseline) is when a wave of depolarization is moving *away* from the positive electrode (towards the negative end of the lead).
- EKG paper has 1 millimeter small squares – so height and depth of wave is measured in millimeters.
10 mm = 1.0 mV
- Horizontal axis is time.
 - 0.04 seconds for 1 mm (1 small box).
 - 0.2 seconds for 1 large box = 5 small boxes = 5 x .04 seconds.

Positive QRS in Lead I.
Negative QRS in Lead aVR.
R wave = 7-8 mm high in Lead I = 7-8mV.
QRS wave = 0.06 seconds long in Lead I.




Rhythm Summary



▶ Rate	90–95 bpm
▶ Regularity	regular
▶ P waves	normal
▶ PR interval	0.12 s
▶ QRS duration	0.08 s
Interpretation?	<i>Normal Sinus Rhythm</i>

Electrocardiogram

- ▶ Intervals show timing of cardiac cycle
 - P-P = one cardiac cycle
 - P-Q = time for atrial depolarization
 - Q-T = time for ventricular depolarization
 - T-P = time for relaxation



- What these intervals represents ?
 - • P wave
 - • PR interval –
 - • QRS Complex –
 - • ST Segment –
 - • T Wave –
 - • U Wave –
 - • RR Interval –




- Intervals
 - P wave – atrial depolarization
 - PR interval – time from sinoatrial node (S-A) to atrioventricular node (A-Vnode)
 - QRS Complex – ventricular depolarization
 - ST Segment – beginning of ventricular repolarization
 - T Wave – later stages of ventricular repolarization
 - U Wave – final component of ventricular repolarization
- RR Interval – represents the time for one complete cardiac cycle



The walls of the left ventricle and the leads that view these walls

The four walls of the left ventricle and the ECG leads that "view" these walls



Anatomic Groups (Septum)

I Lateral	aVR None	V ₁ Septal	V ₄ Anterior
II Inferior	aVL Lateral	V ₂ Septal	V ₅ Lateral
III Inferior	aVF Inferior	V ₃ Anterior	V ₆ Lateral

Anatomic Groups (Anterior Wall)

I Lateral	aVR None	V ₁ Septal	V ₄ Anterior
II Inferior	aVL Lateral	V ₂ Septal	V ₅ Lateral
III Inferior	aVF Inferior	V ₃ Anterior	V ₆ Lateral

Anatomic Groups (Lateral Wall)

I Lateral	aVR None	V ₁ Septal	V ₄ Anterior
II Inferior	aVL Lateral	V ₂ Septal	V ₅ Lateral
III Inferior	aVF Inferior	V ₃ Anterior	V ₆ Lateral


Anatomic Groups (Inferior Wall)

I Lateral	aVR None	V ₁ Septal	V ₄ Anterior
II Inferior	aVL Lateral	V ₂ Septal	V ₅ Lateral
III Inferior	aVF Inferior	V ₃ Anterior	V ₆ Lateral


Anatomic Groups (Summary)


I Lateral	aVR None	V ₁ Septal	V ₄ Anterior
II Inferior	aVL Lateral	V ₂ Septal	V ₅ Lateral
III Inferior	aVF Inferior	V ₃ Anterior	V ₆ Lateral

Which Leads Look Where?



Inferior wall: II, III, aVF
 Septum: V1, V2
 Anterior wall: V3, V4
 Lateral wall: I, aVL (high lateral); V5, V6 (low lateral)
 Anterior wall and septum often infarct together because both supplied by LAD, so anteroseptal MI shows in V1-V4






Coronary Anatomy & ECG Leads

Lateral Leads	I, aVL, V5 - V6	LCx or Diagonal of LAD
Inferior Leads	II, III, aVF	RCA and/or LCx
Anterior/Septal Leads	V1 - V4	LAD



**REBEL
REVIEWS**

Areas of the ECG to be concentrated upon to study the events e.g. MI



Position	Leads
Lateral	look on lead I, V5 ,V6
Inferior	look on lead II ,III ,aVF
Anterior/ Septal	look on V1,V2,V3,V4

Rule of 300

Take the number of “big boxes” between neighboring QRS complexes, and divide this into 300. The result will be approximately equal to the rate

Although fast, this method only works for regular rhythms.






Fig. 1.23 12-lead ECG: example 1


Description

- Sinus rhythm, rate 110/min
- Normal PR interval (140 ms)
- Normal QRS duration (120 ms)
- Normal cardiac axis
- Normal QRS complexes
- Normal T waves (an inverted T wave in lead VR is normal)

Interpretation

- Normal ECG


What is the heart rate?



www.uptodate.com

$$(300 / 6) = 50 \text{ bpm}$$


What is the heart rate?



www.uptodate.com

$$(300 / \sim 4) = \sim 75 \text{ bpm}$$

What is the heart rate?



$$(300 / 1.5) = 200 \text{ bpm}$$

The Rule of 300

It may be easiest to memorize the following table:


# of big boxes	Rate
1	300
2	150
3	100
4	75
5	60
6	50

10 Second Rule

As most EKGs record 10 seconds of rhythm per page, one can simply count the number of beats present on the EKG and multiply by 6 to get the number of beats per 60 seconds.


This method works well for irregular rhythms.





What is the heart rate?

$33 \times 6 = 198 \text{ bpm}$



ECGs, Normal and Abnormal




(a) Sinus rhythm (normal)




(b) Nodal rhythm – no SA node activity

ECGs, Abnormal



Extrasystole : note inverted QRS complex, misshapen QRS and T and absence of a P wave preceding this contraction.

ECGs, Abnormal



(c) Heart block

Arrhythmia: conduction failure at AV node





(e) Ventricular fibrillation



No pumping action occurs

© The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Irregular ECGs




A



B

© The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

Irregular ECGs



C



D

Rate Rhythm Axis **Intervals** Hypertrophy Infarct

PR interval



< 0.12 s	0.12-0.20 s	> 0.20 s
High catecholamine states Wolff-Parkinson-White	Normal	AV nodal blocks



Rate Rhythm Axis **Intervals** Hypertrophy Infarct

QRS complex

$\leq 0.10 \text{ s}$	0.10-0.12 s	$> 0.12 \text{ s}$
Normal	Incomplete bundle branch block	Bundle branch block PVC Ventricular rhythm




Rate Rhythm Axis **Intervals** Hypertrophy Infarct

QT interval

The duration of the QT interval is proportionate to the heart rate.

The faster the heart beats, the faster the ventricles repolarize **so the shorter the QT interval.** Therefore what is a “normal” QT varies with the heart rate. For each heart rate you need to calculate an adjusted QT interval, called the “corrected QT” (QTc):



$$QTc = QT / \text{square root of RR interval}$$




Rate Rhythm Axis **Intervals** Hypertrophy Infarct

QTc interval

$< 0.44 \text{ s}$	$> 0.44 \text{ s}$
Normal	Long QT



Rate Rhythm Axis **Intervals** Hypertrophy Infarct




PR
interval? *0.16
seconds*

QRS
width? *0.08
seconds*


QTc
interval? *0.49
seconds*

Interpretation of
intervals? *Normal PR and QRS, long
QT*

Irregular ECGs





A




B

©The McGraw-Hill Companies, Inc. Permission required for reproduction or display.


Irregular ECGs



C




D




PR = 0.16 s
Normal complex




PR = 0.38 s
First-degree heart block




Second-degree heart block
(2:1 heart block)



Second-degree heart block
(Wenckebach phenomenon)



Complete heart block. Atrial rate, 107; ventricular rate, 43



Two V leads in left
bundle branch block

Figure 28-11. Heart block.




Figure 28-13. Atrial arrhythmias. The illustration shows an atrial premature beat with its P wave superimposed on the T wave of the preceding beat (arrow); atrial tachycardia; atrial flutter with 4:1 AV block; and atrial fibrillation with a totally irregular ventricular rate. (Tracings reproduced, with permission, from Goldschlager N, Goldman MJ: *Principles of Clinical Electrocardiography*, 13th ed. Originally published by Appleton & Lange. Copyright © 1989 by The McGraw-Hill Companies, Inc.)





Figure 28-14. Top: Ventricular premature beats (VPB). The lines under the tracing illustrate the compensatory pause and show that the duration of the premature beat plus the preceding normal beat is equal to the duration of two normal beats. **Bottom:** Ventricular tachycardia.

SINUS node is the pacemaker, firing at a regular rate of 60–100 times per minute. Each beat is conducted normally through to the ventricles.




Normal Sinus Rhythm

SINUS node is the pacemaker, firing regularly at a rate of less than 60 times per minute. Each impulse is conducted normally through to the ventricles.



Sinus Bradycardia


SINUS node is the pacemaker, firing regularly at a rate of greater than 100 times per minute. Each impulse is conducted normally through to the ventricles.



*

Sinus Tachycardia

SINUS node is the pacemaker, but impulses are initiated in an irregular pattern. The rate increases as the patient breathes in and decreases as the patient breathes out. Each beat is conducted normally through to the ventricles.



Sinus Arrhythmia

The pacemaker is an irritable focus within the ATRIUM that fires prematurely and produces a single ectopic beat. Conduction through to the ventricles is normal.



Premature atrial contraction acronym is PAC; sort of a misnomer as this is a premature atrial depolarization.

Some also call this a premature atrial beat (PAB), but again electrical depolarization does not always mean mechanical contraction.

Here there is a focus other than the SA node that is firing, which causes a QRS.


What is the underlying rate? 7×10 using 6-second. Or, 75 using 4 big boxes per QRS.

The PAC is the third QRS; you can see there is only about 1 big box between the 2nd and 3rd QRS.

Usually the abnormal focus will cause a p-wave that looks different from the other p-waves (the SA node p-waves).


Here the aberrant atrial depolarization appears as a peaked p-wave in comparison.


Premature Atrial Contraction



EMEDU


Sinus Tachycardia: rate= 155





EMEDU

Sinus Tachycardia rate: 150 Incomplete LBBB



What do you think?



A: normal
B: sinus tachycardia

a




b




What do you think?



a





b




a A: normal
 B: sinus bradycardia








Normal




Atrial Flutter








Normal



Atrial Fibrillation







Normal



Ventricular Tachycardia (VT)





EMEDU

Atrial Flutter with 2:1 conduction

Mild lateral ST depression V4-V6

aVR

V1

V4

Ventricular Rate= 150

II Flutter waves: Atrial rate =300
buried in the QRS complex

aVL

V2


V5

III

aVF

V3

V6



Atrial Flutter



Heart Blocks

Constant P-R interval

Variable P-R Interval


First Degree Heart Block



Second Degree Heart Block Type I



Second Degree Heart Block Type II



Third Degree Heart Block



First degree AV block



Second degree AV block (Mobitz I or Wenckebach)



Second degree AV block (Mobitz II)




Second degree AV block (2:1 block)




Third degree AV block with junctional escape




First Degree Heart Block



Second degree heart block, type I (Wenckebach/Mobitz I)



Second degree heart block, type 2 (Mobitz 2)



Third degree heart block



SUMMARY

Rate Rhythm Axis Intervals Hypertrophy Infarct

To summarize:

1. Calculate RATE
2. Determine RHYTHM
3. Determine QRS AXIS

-


-

-

-





- ▶ Significant Q wave = Necrosis
- ▶ ST elevation = Injury
- ▶ T wave inversion = Ischemia




Myocardial Infarction

Location of Myocardial Ischemia/ Infarction

Location	Leads
Anterior	I, V ₂ , V ₃ , and V ₄
Anterolateral	I, aVL, V ₅ , and V ₆
Lateral	V ₅ and V ₆
High lateral	I and aVL (often with V ₅ , V ₆)
Inferior	II, III, and aVF
Inferolateral	II, III, aVF, and V ₆
True posterior	Reciprocal changes in V ₁ and V ₂



MI Location



Myocardial involvement

Anterior

Anteroseptal

Anterolateral

Extensive Anterior

V1 through V6 (all)

Lateral

High lateral

Inferior

Inferolateral as above,

Posterior

changes)

Inferoposterolateral

Right Ventricular

V5R

EKG leads

V2, V3, V4 (at least 2)

V1, V2, V3 (+V4)

V4, V5, V6 (+V3, +V2)

V5, V6 (+I, +aVL)

I, aVL

II, III, aVF (at least

2)

+V6 (+V5)

V1, V2 (*recip.

changes)

Combine above 3 items

V4R, +V3R and/or


V5R

SUMMARY

Rate Rhythm Axis Intervals Hypertrophy **Infarct**

Infarct: Is the ST elevation or depression?

*Yes! Elevation in V2-V6, I and avL.
Depression in II, III and avF.*



► ELEVATION

- Electrolytes
- Left bundle branch block
- Early repolarization
- Ventricular hypertrophy
- Aneurysm
- Treatment (pericardiocentesis)
- Injury (acute MI, contusion)
- Osborne waves (hypothermia)
- Nonocclusive vasospasm

ST Segment Elevation




- ▶ **DEPRESSED ST**
 - Drooping valve (mitral valve prolapse)
 - Enlargement of LV with strain
 - Potassium loss (hypokalemia)
 - Reciprocal ST depression (inferior MI)
 - Embolism (PE)
 - Subendocardial ischemia
 - Subendocardial infarct
 - Encephalon hemorrhage
 - Dilated cardiomyopathy
 - Shock
 - Toxicity of digitalis, quinidine

ST Segment Depression




Hyperkalemia

- High K^+
- Peaked T




moderate




extreme

Hypokalemia

- Low K^+
- Flat T, U Wave



moderate



extreme

hyperkalemia



normal P waves




P waves flatten, prolonged PR



P waves disappear, bradycardia

▶ HYPERCALCEMIA




HYPERcalcaemia

▶ HYPOCALCEMIA




Normal




HYPOcalcaemia

Hypercalcemia




Normal QT Interval



Shortened
QT Interval

Causes of Hypercalcemia

- Malignancy
- Milk-alkali syndrome
- Vitamin D toxicity
- Paget's disease
- Hyperparathyroidism
- Immobilization
- Granulomatous disease
- Thiazide diuretics




pyright (your organization) 2003 9/23/2020



Normal tracing (plasma K^+ 4–5.5 meq/L). PR interval = 0.16 s; QRS interval = 0.06 s; QT interval = 0.4 s (normal for an assumed heart rate of 60).



Hyperkalemia (plasma K^+ \pm 7.0 meq/L). The PR and QRS intervals are within normal limits. Very tall, slender peaked T waves are now present.



Hyperkalemia (plasma K^+ \pm 8.5 meq/L). There is no evidence of atrial activity; the QRS complex is broad and slurred and the QRS interval has widened to 0.2 s. The T waves remain tall and slender. Further elevation of the plasma K^+ level may result in ventricular tachycardia and ventricular fibrillation.



Hypokalemia (plasma K^+ \pm 3.5 meq/L). PR interval = 0.2 s; QRS interval = 0.06 s; ST segment depression. A prominent U wave is now present immediately following the T. The actual QT interval remains 0.4 s. If the U wave is erroneously considered a part of the T, a falsely prolonged QT interval of 0.6 s will be measured.




Hypokalemia (plasma K^+ \pm 2.5 meq/L). The PR interval is lengthened to 0.32 s; the ST segment is depressed; the T wave is inverted; a prominent U wave is seen. The true QT interval remains normal.

Figure 28-20. Correlation of plasma K^+ level and the ECG, assuming that the plasma Ca^{2+} level is normal. The diagrammed complexes are left ventricular epicardial leads. (Reproduced, with permission, from Goldman MJ: *Principles of Clinical Electrocardiography*, 12th ed. Originally published by Appleton & Lange. Copyright © 1989 by The McGraw-Hill Companies, Inc.)

ECGs, Normal and Abnormal




(a) Sinus rhythm (normal)




(b) Nodal rhythm – no SA node activity

ECGs, Abnormal



Extrasystole : note inverted QRS complex, misshapen QRS and T and absence of a P wave preceding this contraction.

ECGs, Abnormal



(c) Heart block

Arrhythmia: conduction failure at AV node



(e) Ventricular fibrillation

No pumping action occurs

THE ABNORMAL

Rate Rhythm Axis Intervals **Hypertrophy** Infarct

In this step of the 12-lead ECG analysis, we use the ECG to determine if any of the 4 chambers of the heart are enlarged or hypertrophied. We want to determine if there are any of the following:


- Right atrial enlargement (RAE)
- Left atrial enlargement (LAE)
- Right ventricular hypertrophy (RVH)
- Left ventricular hypertrophy (LVH)



Rate Rhythm Axis Intervals **Hypertrophy** Infarct

Right atrial enlargement

- Take a look at this ECG. What do you notice about the P waves?




***The P waves are tall, especially in leads II, III and aVF.
Ouch! They would hurt to sit on!!***

Rate Rhythm Axis Intervals **Hypertrophy** Infarct

Right atrial enlargement

- To diagnose RAE you can use the following criteria:
 - II $P > 2.5 \text{ mm}$, or
 - V1 or V2P $> 1.5 \text{ mm}$




A cause of RAE is RVH from pulmonary hypertension.

Rate Rhythm Axis Intervals **Hypertrophy** Infarct

Left atrial enlargement

- Take a look at this ECG. What do you notice about the P waves?





The P waves in lead II are notched and in lead V1 they have a deep and wide negative component.

Rate Rhythm Axis Intervals **Hypertrophy** Infarct

Left atrial enlargement

- To diagnose LAE you can use the following criteria:
 - II > 0.04 s (1 box) between notched peaks, or
 - V1 Neg. deflection > 1 box wide x 1 box deep




A common cause of LAE is LVH from hypertension.

Rate Rhythm Axis Intervals **Hypertrophy** Infarct

Right ventricular hypertrophy

- Take a look at this ECG. What do you notice about the axis and QRS complexes over the right ventricle (V1, V2)?





There is right axis deviation (negative in I, positive in II) and there are tall R waves in V1, V2.

Rate Rhythm Axis Intervals **Hypertrophy** Infarct

Right ventricular hypertrophy


- Compare the R waves in V1, V2 from a normal ECG and one from a person with RVH.
- Notice the R wave is normally small in V1, V2 because the right ventricle does not have a lot of muscle mass.
- But in the hypertrophied right ventricle the R wave is tall in V1, V2



Rate Rhythm Axis Intervals **Hypertrophy** Infarct

Right ventricular hypertrophy

- To diagnose RVH you can use the following criteria:
 - Right axis deviation, and
 - V1 R wave > 7mm tall




A common cause of RVH is left heart failure.

Rate Rhythm Axis Intervals **Hypertrophy** Infarct

Left ventricular hypertrophy

- Take a look at this ECG. What do you notice about the axis and QRS complexes over the left ventricle (V5, V6) and right ventricle (V1, V2)?




There is left axis deviation (positive in I, negative in II) and there are tall R waves in V5, V6 and deep S waves in V1, V2

Rate Rhythm Axis Intervals **Hypertrophy** Infarct

Left ventricular hypertrophy


- To diagnose LVH you can use the following criteria :
 - $R \text{ in } V5 \text{ (or } V6) + S \text{ in } V1 \text{ (or } V2) > 35 \text{ mm, or}$
 - avL $R > 13 \text{ mm}$



A common cause of LVH is hypertension.

Rate Rhythm Axis Intervals **Hypertrophy** Infarct

A 63 yo man has longstanding, uncontrolled hypertension. Is there evidence of heart disease from his hypertension? (*Hint: There are 3 abnormalities.*)



Yes, there is left axis deviation (positive in I, negative in II), left atrial enlargement ($> 1 \times 1$ boxes in V1) and LVH (R in V5 = 27 + S in V2 = 10 \rightarrow > 35 mm).

Rate Rhythm Axis Intervals Hypertrophy **Infarct**

- ▶ When analyzing a 12-lead ECG for evidence of an infarction you want to look for the following:
 - **Abnormal Q waves**
 - **ST elevation or depression**
 - **Peaked, flat or inverted T waves**
- ▶ These topics were covered in **Modules V and VI** where you learned:
 - **ST elevation (or depression) of 1 mm in 2 or more contiguous leads is consistent with an AMI**
 - **There are ST elevation (Q-wave) and non-ST elevation (non-Q wave) MIs**

